

2-14-00

A

02/10/00

UTILITY PATENT APPLICATION TRANSMITTAL <small>(Only for new nonprovisional applications under 37 CFR 1.53(b))</small>	<i>Title of Invention</i>	Deglycosylated Kringle 1-5 Region Fragments of Plasminogen and Methods of Use
	<i>Named Inventor(s)</i>	Steven Pirie-Shepherd, Oliver Kisker, M. Judah Folkman, Kim Lee Sim, Nicholas MacDonald and Hong Li Ang
	<i>Attorney Docket</i>	05940-0141
	<i>Express Mail Label No.</i>	EL498678449US

02/10/00
PTO
91705676
1065 U.S.

APPLICATION ELEMENTS	Assistant Commissioner for Patents ADDRESS TO: Box Patent Application Washington, D.C. 20231
1. <input type="checkbox"/> Fee Transmittal Form <small>(Submit an original, and a duplicate for fee processing)</small> 2. <input checked="" type="checkbox"/> Specification, Claims, and Abstract Total Pages 43 3. <input checked="" type="checkbox"/> Drawings Total Sheets 4 4. Oath or Declaration Total Pages _____ a. <input type="checkbox"/> Newly executed (original or copy) b. <input type="checkbox"/> Copy from prior application (37 CFR 1.63(d)) <small>(for continuation/divisional with Box 17 completed)</small> [Note Box 5 Below] (i) <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b). 5. <input type="checkbox"/> Incorporation by Reference <small>(usable if Box 4b is checked)</small> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein. 6. <input type="checkbox"/> Microfiche Computer Program <i>(Appendix)</i> 7. <input type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission <i>(if applicable, all necessary)</i> a. <input type="checkbox"/> Computer Readable Copy b. <input type="checkbox"/> Paper Copy (identical to computer copy) c. <input type="checkbox"/> Statement verifying identity of above copies	ACCOMPANYING APPLICATION PARTS 8. <input type="checkbox"/> Assignment Papers (cover sheet & document(s)) 9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <small>(when there is an assignee)</small> <input type="checkbox"/> Power of Attorney by assignee 10. <input type="checkbox"/> English Translation Document <i>(if applicable)</i> 11. <input type="checkbox"/> Information Disclosure Statement (IDS) PTO-1449 <input type="checkbox"/> Copies of IDS Citations 12. <input type="checkbox"/> Preliminary Amendment 13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) <small>(Should be specifically itemized)</small> 14. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application Status still proper and desired 15. <input type="checkbox"/> Certified Copy of Priority Document(s) 16. <input type="checkbox"/> Other: _____ _____ _____
17. If a CONTINUING APPLICATION , check appropriate box and supply the requisite information: <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application No: _____	
18. CORRESPONDENCE ADDRESS: <div style="display: flex; justify-content: space-between;"> <div> Lisa C. Elsevier, Esq. JONES & ASKEW, LLP 2400 Monarch Tower 3424 Peachtree Road, N.E. Atlanta, Georgia 30326 </div> <div> By: <u>Lisa C. Elsevier</u> Reg. No. 44.669 Date: <u>February 10, 2000</u> Telephone: 404-949-2400 Facsimile: 404-949-2499 </div> </div>	

5

**DEGLYCOSYLATED KRINGLE 1-5 REGION
FRAGMENTS OF PLASMINOGEN
AND METHODS OF USE**

10

This invention was made with government support under Grant No.(s) CA 45548 by the NIH. The government may have certain rights in the invention.

15

Cross Reference to Related Applications

This application claims priority to U.S. Provisional Patent Application Serial No. 60/119,562, filed February 10, 1999 and U.S. Provisional Patent Application Serial No. 60/128,062, filed April 7, 1999.

20

Field of the Invention

The present invention relates to compositions and methods for the modulation of angiogenesis. More particularly, the present invention relates to deglycosylated kringle 1-5 region proteins that are useful for the treatment of angiogenesis-associated diseases such as cancer.

25

30

Background of the Invention

As used herein, the term "angiogenesis" means the generation of new blood vessels into a tissue or organ. Under normal physiological conditions, humans or animals undergo angiogenesis only in very specific restricted situations. For example, angiogenesis is normally observed in wound healing.

35

fetal and embryonal development and formation of the corpus luteum, endometrium and placenta. However, angiogenesis also occurs under abnormal or undesired conditions such as during tumor development, growth and metastasis. This type of angiogenesis may also be referred to as uncontrolled angiogenesis.

Both controlled and uncontrolled angiogenesis are thought to proceed in a similar manner. Endothelial cells and pericytes surrounded by a basement membrane form capillary blood vessels. Angiogenesis begins with the erosion of the basement membrane by enzymes released by endothelial cells and leukocytes. The endothelial cells, which line the lumen of blood vessels, then protrude through the basement membrane. Angiogenic stimulants induce the endothelial cells to migrate through the eroded basement membrane. The migrating cells form a "sprout" off the parent blood vessel, where the endothelial cells undergo mitosis and proliferate. The endothelial sprouts merge with each other to form capillary loops, creating the new blood vessel.

Persistent, unregulated angiogenesis occurs in a multiplicity of disease states, tumor metastasis and abnormal growth by endothelial cells. The diverse pathological disease states in which unregulated angiogenesis is present have been grouped together as angiogenic dependent or angiogenic associated diseases. The hypothesis that tumor growth is angiogenesis-dependent was first proposed in 1971 by M. Judah Folkman. (Folkman J., "Tumor angiogenesis: Therapeutic implications" *N. Engl. Jour. Med.* 285:1182-1186 (1971)). In its simplest terms the hypothesis states: "Once tumor 'take' has occurred, every increase in tumor cell population must be preceded by an increase in new capillaries converging on the tumor." Tumor 'take' is currently understood to indicate a pre-vascular phase of tumor growth in which a population of tumor cells occupying a few cubic millimeters volume and not exceeding a few million cells, can survive on existing host

microvessels. Expansion of tumor volume beyond this phase requires the induction of new capillary blood vessels.

Several molecules have been discovered that inhibit angiogenesis, inhibit tumor growth, cause regression of primary tumors, and/or inhibit metastasis of primary tumors. These molecules are called antiangiogenic agents. One of these antiangiogenic agents has been termed angiostatin, which is a fragment of a plasminogen protein. Angiostatin was first described in U.S. Patent No. 5,639,725.

Plasminogen protein comprises five kringle region domains and a serine protease domain located at the carboxy-terminal region. The DNA sequence of human plasminogen has been published. (Browne, M.J., *et al.*, "Expression of recombinant human plasminogen and aglycoplasminogen in HeLa cells" *Fibrinolysis* 5(4):257-260 (1991)). Each kringle region of the plasminogen molecule contains approximately 80 amino acids and contains 3 disulfide bonds. (Robbins, K.C., "The plasminogen-plasmin enzyme system" *Hemostasis and Thrombosis, Basic Principles and Practice*, 2nd Edition, ed. by Colman, R.W. *et al.* J.B. Lippincott Company, pp. 340-357, 1987). The approximate amino acid spans of each kringle domain of human plasminogen are as follows: kringle 1 typically encompasses Cys84-Cys162, kringle 2 typically encompasses Cys166-Cys243, kringle 3 typically encompasses Cys256-Cys333, kringle 4 typically encompasses Cys358-Cys435, and kringle 5 typically encompasses Cys462-Cys541. (Castellino, F.J. and McCance, S.G., "The kringle domains of human plasminogen" *Ciba Found. Symp.*, 212:46-65 (1997)). Because each kringle domain is separated by an approximately 20 amino acid inter-kringle domain, a kringle region as herein defined can include any portion of these adjacent inter-kringle domains.

Angiostatin protein comprises one or more of these five kringle regions of plasminogen. Among the proteases suggested to be responsible for angiostatin generation are macrophage metalloelastase, metalloproteinases (mmp's) -3, - 7, and -9 and

plasmin itself in the presence of a free sulphhydryl donor such as cysteine. (Dong, Z. *et al.* "Macrophage-derived metalloelastase is responsible for the generation of angiostatin in Lewis lung carcinoma" *Cell*, 88:801-810 (1997); Lijnen, H.R. *et al.*, "Generation of an angiostatin-like fragment from plasminogen by stromelysin-1 (MMP-3)" *Biochemistry*, 37:4699-4702 (1998); Patterson, B.C. and Sang Q.A., "Angiostatin-converting enzyme activities of human matrilysin (MMP-7) and gelatinase B/type IV collagenase (MMP-9)" *J. Biol. Chem.* 272:28823-28825 (1997)) (Stathakis, P. *et al.*, "Generation of angiostatin by reduction and proteolysis of plasmin" *J. Biol. Chem.* 272:20641-20645 (1997); Gately, S. *et al.*, "The mechanism of cancer-mediated conversion of plasminogen to that angiogenesis inhibitor angiostatin" *Proc. Natl. Acad. Sci. USA* 94:10868-10872 (1997)).

In its glycosylated state, human plasminogen protein contains an N-linked carbohydrate chain at amino acid position Asn-289 and two O-linked mucin type carbohydrate chains at amino acid positions Ser-249 and Thr-346. Since angiostatin protein is a fragment of a plasminogen protein, angiostatin protein also contains the above-described carbohydrate chains. The proteolytic digestion of plasminogen, by tPA and uPA to produce plasmin for example, is known to be modulated by the carbohydrate content of plasminogen. Further the carbohydrate chains are known to modulate binding of plasminogen to cell surface receptors. Carbohydrate is also known to play a general role in systemic half life of circulating proteins. (Mori, K. *et al.*, "The activation of type 1 and type 2 plasminogen by type I and type II tissue plasminogen activator" *J. Biol. Chem.* 270:3261-3267 (1995); Davidson, D.J. and Castellino, F.J., "The influence of the nature of the asparagine 289-linked oligosaccharide on the activation by urokinase and lysine binding properties of natural and recombinant human plasminogens" *J. Clin. Invest.* 92:249-254 (1993); Edelberg, J. *et al.* "Neonatal plasminogen displays altered cell surface binding and activation kinetics" *J. Clin.*

Invest. 86:107-112 (1990); Gonzales-Gronow, M. *et al.* "Further characterization of the cellular plasminogen binding site: evidence that plasminogen 2 and lipoprotein a compete for the same site" *Biochemistry*, 28:2374-2377 (1989); Jenkins, N. *et al.* "Getting the glycosylation right: implications for the biotechnology industry" *Nat. Biotechnol.* 14:975-981 (1996)).

The mechanism underlying how angiostatin and its related kringle fragments specifically inhibit endothelial cell growth remains uncharacterized. It is not yet clear whether the inhibition is mediated by a receptor that is specifically expressed in proliferating endothelial cells, or if angiostatin is internalized by endothelial cells and subsequently inhibits cell proliferation. Alternatively, angiostatin may interact with an endothelial cell adhesion receptor such as integrin $\alpha_v\beta_3$, blocking integrin-mediated angiogenesis (Brooks, P.C., *et al.* "Integrin α_v β_3 antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels" *Cell* 79:1157-1164 (1994)).

Although angiostatin has been identified as an angiogenesis inhibitor, what is needed in the art are kringle region fragments of plasminogen that have increased antiangiogenic activity. These improved kringle region proteins should be useful for the treatment of angiogenesis-mediated diseases, such as cancer, and for the modulation of other angiogenic processes, such as wound healing and reproduction. Due to the improved nature of the antiangiogenic kringle region proteins, these proteins should be able to be administered in smaller doses, thus lowering the cost of cancer treatment. What is also needed in the art are compositions and methods for the detection, measurement and localization of improved antiangiogenic kringle region proteins.

Summary of the Invention

Compositions and methods are provided herein that are effective for modulating angiogenesis, and inhibiting unwanted

angiogenesis, especially angiogenesis related to tumor growth. In particular, the present invention provides deglycosylated kringle 1-5 region proteins that are fragments of plasminogen. It is a surprising discovery of the present invention that deglycosylation of kringle 1-5 region proteins dramatically increases the antiangiogenic activity of these proteins. It is to be understood that the deglycosylated kringle 1-5 region proteins described herein include fusion proteins wherein a deglycosylated kringle 1-5 region protein is contiguous with one or more other proteins. In a preferred embodiment, the fusion protein comprises a deglycosylated kringle 1-5 region protein and another antiangiogenic protein.

Also included in the present invention are nucleotides encoding the deglycosylated kringle 1-5 region proteins, expression vectors containing DNA sequences encoding deglycosylated kringle 1-5 region proteins, and cells containing one or more expression vectors containing DNA sequences encoding deglycosylated kringle 1-5 region proteins.

The present invention also includes antibodies specific for the deglycosylated kringle 1-5 region proteins, antibodies that inhibit the binding of antibodies specific for deglycosylated kringle 1-5 region proteins, and antibodies specific for a deglycosylated kringle 1-5 region protein receptor. These antibodies can be polyclonal antibodies or monoclonal antibodies. The antibodies specific for the deglycosylated kringle 1-5 region proteins can be used in diagnostic kits to detect the presence and quantity of deglycosylated kringle 1-5 region proteins which is diagnostic or prognostic for the occurrence or recurrence of cancer or other diseases mediated by angiogenesis. Antibodies specific for deglycosylated kringle 1-5 region proteins may also be administered to a human or animal to passively immunize the human or animal against deglycosylated kringle 1-5 region proteins, thereby reducing angiogenic inhibition.

The present invention provides methods and compositions for treating diseases and processes mediated by undesired and

uncontrolled angiogenesis by increasing an *in vivo* concentration of deglycosylated kringle 1-5 region protein in a human or animal. *In vivo* concentrations of deglycosylated kringle 1-5 region proteins may be increased by administering to a human or animal a composition comprising a substantially purified deglycosylated kringle 1-5 region protein in a dosage sufficient to inhibit angiogenesis. Additionally, *in vivo* concentrations of deglycosylated kringle 1-5 region proteins may be increased in a human or animal by the administration of nucleotides encoding deglycosylated kringle 1-5 region proteins or enzymes that release deglycosylated kringle 1-5 region proteins from plasminogen or plasmin. The present invention is particularly useful for treating or repressing the growth of tumors.

Accordingly, it is an object of the present invention to provide a composition comprising a deglycosylated kringle 1-5 region protein.

It is another object of the present invention to provide a nucleotide composition encoding a deglycosylated kringle 1-5 region protein.

It is another object of the present invention to provide compositions and methods for increasing an *in vivo* concentration of deglycosylated kringle 1-5 region proteins.

It is another object of the present invention to provide a composition comprising an antibody to a deglycosylated kringle 1-5 region protein that is specific for the deglycosylated kringle 1-5 region protein and does not recognize plasminogen.

It is an object of present invention to provide a method for detecting and quantifying the presence of an antibody specific for a deglycosylated kringle 1-5 region protein in a body fluid.

It is another object of the present invention to provide a method for the detection or prognosis of cancer.

It is another object of the present invention to provide a method of treating diseases and processes that are mediated by angiogenesis.

It is another object of the present invention to provide a composition for treating or repressing the growth of a cancer.

Yet another object of the invention is to provide compositions and methods useful for gene therapy for the modulation of angiogenic processes.

These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiments and the appended claims.

Brief Description of the Figures

Figure 1 shows the amino acid sequence (SEQ ID NO:1) and nucleotide sequence (SEQ ID NO:2) encoding a preferred deglycosylated kringle 1-5 region protein.

Figure 2 shows the purification of glycosylated and deglycosylated kringle 1-5 region proteins using lys-Sepharose.

Figure 3 shows the purification of glycosylated and deglycosylated kringle 1-5 region proteins using gel filtration chromatography.

Figure 4 shows the inhibition of endothelial cell proliferation by glycosylated and deglycosylated kringle 1-5 region proteins.

Detailed Description

The present invention includes deglycosylated fragments of a kringle 1-5 region of a plasminogen protein. Also included in the present invention are methods of using the deglycosylated kringle 1-5 region proteins and methods and compositions for detecting the deglycosylated kringle 1-5 region proteins.

A surprising discovery of the present invention is that deglycosylated fragments of the kringle 1-5 region of plasminogen are dramatically more antiangiogenic than

glycosylated fragments of the kringle 1-5 region of plasminogen. Accordingly, the present invention includes deglycosylated fragments of kringle 1-5 region proteins and compositions comprising deglycosylated fragments of kringle 1-5 region proteins. In one embodiment, a composition comprises a pharmaceutically acceptable carrier and a deglycosylated fragment of a kringle 1-5 region of a plasminogen protein in a greater concentration than a naturally glycosylated form of the deglycosylated fragment, wherein the deglycosylated fragment lacks one or more carbohydrate moieties linked to the naturally glycosylated form and wherein the deglycosylated fragment has antiangiogenic activity. In a further embodiment, the ratio of the amount of the deglycosylated fragment to the amount of the naturally glycosylated form is at least 60:40, more preferably at least 80:20, and most preferably 100:0. The ratio of the deglycosylated fragment to the naturally glycosylated form may also be at least 95:5, 96:4, 97:3, 98:2 and 99:1.

As used herein, the term “deglycosylated” refers to a protein or peptide lacking one or more carbohydrate moieties that are linked to a naturally glycosylated form of a kringle 1-5 region proteins. A “naturally glycosylated form” refers to a fragment of a kringle 1-5 region protein, or a larger protein from which the kringle 1-5 region fragment is derived, that is glycosylated at amino acid positions as found in nature. For example, one naturally glycosylated form of human plasminogen is glycosylated at Ser-249, Asn-289, and Thr-346, and therefore, the naturally glycosylated form of fragments of a kringle 1-5 region of human plasminogen are glycosylated at Ser-249, Asn-289, and/or Thr-346. It is to be understood that the naturally glycosylated forms of the kringle 1-5 region fragments described herein are not required to exist as a fragment in nature and can exist in nature as part of a larger protein.

Therefore, “deglycosylated plasminogen” refers to a plasminogen protein of any species that lacks carbohydrate at a position that may be otherwise glycosylated in nature (i.e., that is

glycosylated in a naturally glycosylated form of the plasminogen protein). For example, included in the present invention is a plasminogen protein lacking a bisialylated-biantennary glycan at an amino acid position corresponding to amino acid position 289 of human plasminogen (Asn-289), an O-linked mucin type carbohydrate chain at an amino acid position corresponding to amino acid position 249 of human plasminogen (Ser-249), and/or an O-linked mucin type carbohydrate chain at an amino acid position corresponding to amino acid position 346 of human plasminogen (Thr-346). Additionally, the term “deglycosylated plasminogen” encompasses a plasminogen protein having zero, one or two carbohydrate chains. In one embodiment of the present invention, a deglycosylated plasminogen lacks any N-linked carbohydrate chain at amino acid position corresponding to amino acid position 289 of human plasminogen. In a further embodiment, a deglycosylated plasminogen lacks any carbohydrate moiety at an amino acid position corresponding to amino acid position 289 of human plasminogen.

As also used herein, the terms “deglycosylated kringle 1-5 region protein” and “deglycosylated fragment” refer to a fragment of a kringle 1-5 region of a plasminogen protein of any species that lacks carbohydrate at a position that may be otherwise glycosylated in nature (i.e., that is glycosylated in a naturally glycosylated form of the deglycosylated kringle 1-5 region protein). For example, included herein are deglycosylated kringle 1-5 region proteins that lack a bisialylated-biantennary glycan at an amino acid position corresponding to amino acid position 289 of human plasminogen, an O-linked mucin type carbohydrate chain at amino acid position corresponding to amino acid position 249 of human plasminogen, and/or an O-linked mucin type carbohydrate chain at amino acid position corresponding to amino acid position 346 of human plasminogen. Therefore, the term “deglycosylated kringle 1-5 region protein” encompasses a kringle 1-5 protein having zero, one or two carbohydrate chains. In one embodiment of the

present invention, the deglycosylated kringle 1-5 region protein lacks a bisialylated-biantennary glycan at amino acid position 289. In another embodiment, the carbohydrate chain at amino acid position 289 is smaller than a bisialylated-biantennary glycan. In another embodiment, the deglycosylated kringle 1-5 region protein lacks a carbohydrate moiety at an amino acid position corresponding to amino acid position 289 of human plasminogen.

“Corresponding to”, when referring to amino acids, indicates the comparison of two amino acids in the same region of different proteins, or fragments thereof, wherein the proteins are homologs, orthologs or paralogs. Homologs are defined as proteins with substantial homology, wherein “substantial homology” is defined below. Orthologs are defined as proteins having non-identical amino acid sequences and similar functional characteristics, wherein the proteins are from different species, but wherein the species have a common ancestral origin. Paralogs are defined as proteins having non-identical amino acid sequences and similar functional characteristics, wherein the proteins are from the same species. In the present invention, an example of an amino acid corresponding to Asn-289 located in kringle 3 of human plasminogen is an amino acid located in kringle 3 of murine plasminogen that can have an N-linked carbohydrate moiety attached thereto, and preferably a bisialylated-biantennary glycan attached thereto, as found in nature. It is to be understood that the present invention includes fragments of kringle 1-5 regions of a plasminogen from a species other than a human.

In another embodiment of the present invention, the kringle 1-5 region protein has a modified amino acid at a position corresponding to position 289 of human plasminogen such that a carbohydrate moiety is not added to that position during post-translational modification of the protein. In a preferred embodiment, the amino acid substitution is a conservative substitution. In a further preferred embodiment,

the amino acid substitution is from asparagine to glutamic acid. In yet a further preferred embodiment, the kringle 1-5 region protein has the amino acid sequence as shown in SEQ ID NO:1 that corresponds to an approximately kringle 1-3 region.

5 The deglycosylated kringle 1-5 region proteins of the present invention can be made *in vivo* or *in vitro*. For example, the deglycosylated kringle 1-5 region proteins can be isolated from body fluids including, but not limited to, serum, urine and ascites, or synthesized by chemical or biological methods (e.g. recombinant gene expression, chemical synthesis of
10 oligonucleotides, and *in vitro* enzymatic catalysis of larger proteins such as plasminogen or plasmin). Recombinant techniques include gene amplification from DNA sources using the polymerase chain reaction (PCR), and gene amplification from RNA sources using reverse transcriptase/PCR. For
15 example, a deglycosylated kringle 1-5 region protein can be recombinantly produced by expressing a gene encoding the protein, wherein an amino acid position that can be glycosylated in nature is substituted for an amino acid that is not glycosylated. This gene may be administered to an individual using the gene
20 therapy methods described in more detail below. The kringle 1-5 region proteins can also be made *in vivo* by the administration of deglycosylated plasminogen and an enzyme that cleaves deglycosylated plasminogen to an individual. In a preferred
25 embodiment, the enzyme is an elastase enzyme such as macrophage metalloelastase. Deglycosylated kringle 1-5 proteins can also be made *in vivo* by the administration of an enzyme that specifically cleaves deglycosylated plasminogen.

30 When isolating deglycosylated kringle 1-5 region proteins from larger proteins such as plasminogen or plasmin, either the plasminogen or plasmin may be deglycosylated prior to enzymatic catalysis or the resulting kringle 1-5 region protein itself may be deglycosylated. Methods for deglycosylation of proteins are well known to those skilled in the art. However, it
35 is a surprising discovery of the present invention that there is an

increased yield of kringle 1-5 region proteins following enzymatic catalysis of deglycosylated plasminogen by elastase as compared to enzymatic catalysis of glycosylated plasminogen by elastase. Therefore, in a preferred embodiment of the present invention, kringle 1-5 region proteins are isolated by elastase digestion of deglycosylated plasminogen.

It is to be understood that "kringle 1-5 region" is defined herein as a region that corresponds to approximately amino acid position 1 through approximately amino acid position 541 of human plasminogen. Thus, the deglycosylated fragments of the kringle 1-5 region of the present invention may contain the N-terminal sequence proceeding the kringle 1 region, kringle regions 1, 2, 3, 4, and 5, inter-kringle regions, and antiangiogenic fragments thereof, wherein the aforementioned regions are found in nature as contiguous sequences or not. Antiangiogenic fragments of the kringle 1-5 region that can be deglycosylated are further disclosed in U.S. Patent Nos. 5,639,725 and 5,837,682.

It is also to be understood that the present invention is contemplated to include any deglycosylated kringle 1-5 region protein derivatives. A kringle 1-5 region protein derivative includes a protein having the amino acid sequence of a kringle 1-5 region of a plasminogen protein. A kringle 1-5 region protein derivative also includes a peptide having a sequence corresponding to an antiangiogenic fragment a kringle 1-5 region. An "antiangiogenic fragment" is defined to be a peptide whose amino acid sequence corresponds to a subsequence of a kringle 1-5 region, referred to as an "antiangiogenic subsequence". A "subsequence" is a sequence of contiguous amino acids found within a larger sequence. A subsequence is generally composed of approximately at least 70%, more preferably 80%, and most preferably 90% of the larger sequence.

A kringle 1-5 region protein derivative also includes a protein or peptide having a modified sequence in which one or

more amino acids in the original sequence or subsequence have been substituted with a naturally occurring amino acid residue or amino acid residue analog (also referred to as modified amino acid). Suitable kringle 1-5 region protein derivatives have

An "amino acid residue" is a moiety found within a protein or peptide and is represented by -NH-CHR-CO- , wherein R is the side chain of a naturally occurring amino acid. When referring to a moiety found within a peptide, the terms "amino acid residue" and "amino acid" are used interchangeably. An "amino acid residue analog" includes D or L configurations having the following formula: -NH-CHR-CO- , wherein R is an aliphatic group, a substituted aliphatic aromatic group, a benzyl group, a substituted benzyl group, an aromatic group or a substituted aromatic group and wherein R does not correspond to the side chain of a naturally occurring amino acid.

Suitable substitutions for amino acid residues in the sequence of the deglycosylated kringle 1-5 region proteins described herein include conservative substitutions that result in angiogenic deglycosylated kringle 1-5 region protein derivatives. A conservative substitution is a substitution in which the substituting amino acid (naturally occurring or modified) is structurally related to the amino acid being substituted. "Structurally related" amino acids are approximately the same size and have the same or similar functional groups in the side chains.

Provided below are groups of naturally occurring and modified amino acids in which each amino acid in a group has similar electronic and steric properties. Thus, a conservative substitution can be made by substituting an amino acid with another amino acid from the same group. It is to be understood that these groups are non-limiting and that additional modified amino acids could be included in each group.

Group I includes leucine, isoleucine, valine, methionine and modified amino acids having the following side chains: ethyl, *n*-propyl *n*-butyl. Preferably, Group I includes leucine, isoleucine, valine and methionine.

5 Group II includes glycine, alanine, valine and a modified amino acid having an ethyl side chain. Preferably, Group II includes glycine and alanine.

10 Group III includes phenylalanine, phenylglycine, tyrosine, tryptophan, cyclohexylmethyl, and modified amino residues having substituted benzyl or phenyl side chains. Preferred substituents include one or more of the following: halogen, methyl, ethyl, nitro, $-NH_2$, methoxy, ethoxy and $-CN$. Preferably, Group III includes phenylalanine, tyrosine and tryptophan.

15 Group IV includes glutamic acid, aspartic acid, a substituted or unsubstituted aliphatic, aromatic or benzylic ester of glutamic or aspartic acid (e.g., methyl, ethyl, *n*-propyl *iso*-propyl, cyclohexyl, benzyl or substituted benzyl), glutamine, asparagine, $-CO-NH$ -alkylated glutamine or asparagine (e.g., methyl, ethyl, *n*-propyl and *iso*-propyl) and modified amino acids having the side chain $-(CH_2)_3-COOH$, an ester thereof (substituted or unsubstituted aliphatic, aromatic or benzylic ester), an amide thereof and a substituted or unsubstituted *N*-alkylated amide thereof. Preferably, Group IV includes glutamic acid, aspartic acid, methyl aspartate, ethyl aspartate, benzyl aspartate and methyl glutamate, ethyl glutamate and benzyl glutamate, glutamine and asparagine.

25 Group V includes histidine, lysine, ornithine, arginine, *N*-nitroarginine, β -cycloarginine, γ -hydroxyarginine, *N*-amidinocitruline and 2-amino-4-guanidinobutanoic acid, homologs of lysine, homologs of arginine and homologs of ornithine. Preferably, Group V includes histidine, lysine, arginine and ornithine. A homolog of an amino acid includes from 1 to about 3 additional or subtracted methylene units in the side chain.

35

Group VI includes serine, threonine, cysteine and modified amino acids having C1-C5 straight or branched alkyl side chains substituted with -OH or -SH, for example, -CH₂CH₂OH, -CH₂CH₂CH₂OH or -CH₂CH₂OHCH₃. Preferably, Group VI includes serine, cysteine or threonine.

In another aspect of the present invention, suitable substitutions for amino acid residues in the amino acid sequences described herein include "severe substitutions" that result in kringle 1-5 region protein derivatives that are antiangiogenic. Severe substitutions that result in antiangiogenic kringle 1-5 region protein derivatives are much more likely to be possible in positions that are not highly conserved than at positions that are highly conserved. In the present invention, severe substitutions are much more likely to be possible in the inter-kringle regions and the N-terminal sequence proceeding kringle 1. A "severe substitution" is a substitution in which the substituting amino acid (naturally occurring or modified) has significantly different size and/or electronic properties compared with the amino acid being substituted. For example, the side chain of the substituting amino acid can be significantly larger (or smaller) than the side chain of the amino acid being substituted and/or can have functional groups with significantly different electronic properties than the amino acid being substituted.

Examples of severe substitutions of this type include the substitution of phenylalanine or cyclohexylmethyl glycine for alanine, isoleucine for glycine, a D amino acid for the corresponding L amino acid or -NH-CH[(-CH₂)₅-COOH]-CO- for aspartic acid. Alternatively, a functional group may be added to the side chain, deleted from the side chain or exchanged with another functional group. Examples of severe substitutions of this type include adding an amine or hydroxyl, carboxylic acid to the aliphatic side chain of valine, leucine or isoleucine, exchanging the carboxylic acid in the side chain of aspartic acid or glutamic acid with an amine or deleting the amine group in the side chain of lysine or ornithine. In yet another alternative,

the side chain of the substituting amino acid can have significantly different steric and electronic properties that the functional group of the amino acid being substituted. Examples of such modifications include tryptophan for glycine, lysine for aspartic acid and $-(CH_2)_4COOH$ for the side chain of serine. These examples are not meant to be limiting.

“Substantial homology” exists between two amino acid sequences when a sufficient number of amino acid residues at corresponding positions of each amino acid sequence are either identical or structurally related such that a protein or peptide having the first amino acid sequence and a protein or peptide having the second amino acid sequence exhibit similar biological activities. Generally, there is substantial sequence homology among the amino acid sequences when at least 30%, more preferably at least 40%, and most preferably at least 50%, of the amino acids in the first amino acid sequence are identical to or structurally related to the second amino acid sequence. Homology is often measured using sequence analysis software, e.g., BLASTIN or BLASTP (available at <http://www.ncbi.nlm.nih.gov/BLAST>). The default parameters for comparing the two sequences (e.g., “Blast”-ing two sequences against each other) by BLASTIN (for nucleotide sequences) are reward for match = 1, penalty for mismatch = -2, open gap = 5, and extension gap = 2. When using BLASTP for protein sequences, the default parameters are reward for match = 0, penalty for mismatch = 0, open gap = 11, and extension gap = 1.

With regard to the present invention, the kringle region motif of kringles 1-3 of a plasminogen protein is also known to exist in other biologically active proteins. These proteins include, but are not limited to, prothrombin, hepatocyte growth factor, scatter factor and macrophage stimulating protein. (Yoshimura, T, *et al.*, “Cloning, sequencing, and expression of human macrophage stimulating protein (MSP, MST1) confirms MSP as a member of the family of kringle proteins and locates the MSP

gene on Chromosome 3" *J. Biol. Chem.*, 268:15461-15468, (1993)). It is contemplated that any kringle 1-5 region protein or peptide having a three-dimensional kringle-like conformation or cysteine motif that has antiangiogenic activity *in vivo*, is part of the present invention.

It is to be understood that as used herein, the term "isolated" refers to a composition which is substantially or essentially free from at least some of the components that normally accompany it in its native state. Thus, the deglycosylated kringle 1-5 region proteins of this invention do not contain some of the materials normally associated with their *in situ* environment. Typically, the isolated, deglycosylated kringle region 1-5 proteins of the invention are at least about 80% pure, usually at least about 90% pure, and preferably at least about 95% pure as measured by band intensity on a silver stained gel. It is to be understood that the term "isolated" does not exclude fusion proteins comprising the deglycosylated kringle 1-5 region proteins from the present invention. The present invention contemplates fusion proteins, or chimeric proteins, comprising the kringle 1-5 region proteins described herein and other proteins. In a preferred embodiment the fusion proteins comprise deglycosylated kringle 1-5 region proteins and other antiangiogenic proteins such as endostatin proteins, wherein the endostatin proteins are defined as antiangiogenic fragments of the C-terminal non-collagenous region of a collagen protein.

The deglycosylated kringle 1-5 region proteins described herein are useful for treating diseases or processes that are mediated by, or involve, angiogenesis. In particular, the deglycosylated kringle 1-5 region proteins are useful for the treatment of angiogenesis-mediated cancers. The deglycosylated kringle 1-5 region proteins are also useful for the generation of antibodies specific for the deglycosylated kringle 1-5 region proteins, and for the identification and isolation of receptors for

and chemical mimetics of deglycosylated kringle 1-5 region proteins.

In addition to deglycosylated kringle 1-5 region proteins, the present invention encompasses compositions comprising nucleotide sequences encoding the kringle 1-5 region proteins described herein. In one embodiment of the present invention, the nucleotide sequence encodes a kringle 1-5 region protein containing a modified amino acid at a position corresponding to amino acid 289 of human plasminogen. In a preferred embodiment, the nucleotide sequence is as shown in SEQ ID NO:2. The present invention also includes a vector containing a DNA sequence encoding a kringle 1-5 region protein, wherein the vector is capable of expressing kringle 1-5 region proteins when present in a cell. The cell may contain one vector or multiple vectors. The nucleotide sequences described herein are useful for recombinantly producing the deglycosylated kringle 1-5 region proteins of the present invention and for treatment of angiogenesis associated diseases and conditions via gene therapy.

Still further, the deglycosylated kringle 1-5 region proteins can be used to generate antibodies to the deglycosylated kringle 1-5 region proteins and their receptors. In particular, an antibody can be generated that specifically binds to a deglycosylated kringle 1-5 region but does not bind to glycosylated plasminogen or a glycosylated kringle 1-5 region protein. The terms "antibody" and "antibodies" as used herein include monoclonal, polyclonal, chimeric, single chain, bispecific, simianized, and humanized antibodies as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

To enhance the potential for high specificity in the development of antisera (or agonists and antagonists) to deglycosylated kringle 1-5 region proteins, protein sequences can be compared to known sequences using protein sequence databases such as GenBank, Brookhaven Protein, SWISS-PROT, and PIR to determine potential sequence homologies. This

information facilitates elimination of sequences that exhibit a high degree of sequence homology to other molecules. These antibodies that specifically bind to the deglycosylated kringle 1-5 region proteins or their receptors, can be used in diagnostic methods and kits that are well known to those of ordinary skill in the art to detect or quantify the deglycosylated kringle 1-5 region proteins or their receptors in a body fluid or tissue. Results from these tests can be used to diagnose or predict the occurrence or recurrence of a cancer or other angiogenic-mediated disease.

The phrases "specifically binds to" or "specific for" when referring to an antibody refers to a binding reaction which is determinative of the presence of the peptide in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind preferentially to a particular peptide and do not bind in a significant amount to other proteins present in the sample. Specific binding to a peptide under such conditions requires an antibody that is selected for its specificity for a particular protein. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See, Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

When labeled with a detectable biomolecule or chemical, the deglycosylated kringle 1-5 region proteins, nucleic acids and antibodies described above are useful for purposes such as *in vivo* and *in vitro* diagnostics and laboratory research using the methods and assays described below. Various types of labels and methods of conjugating the labels to the polypeptides and antibodies are well known to those skilled in the art. Examples

of labels include, but are not limited to, radiolabels, bioluminescent labels, fluorogens and chromogens. Still further, the present invention encompasses deglycosylated kringle 1-5 region proteins, deglycosylated kringle 1-5 region protein antisera, deglycosylated kringle 1-5 region protein receptor agonists or deglycosylated kringle 1-5 region protein receptor antagonists that are combined with pharmaceutically acceptable excipients, and optionally sustained-release compounds or compositions, such as biodegradable polymers, to form therapeutic compositions for use in the methods of treatment described below.

The methods of the present invention include methods of making, detecting and measuring the deglycosylated kringle 1-5 region proteins of the present invention as well as methods of treating angiogenesis associated conditions and diseases using the deglycosylated kringle 1-5 region proteins of the present invention. Deglycosylated kringle 1-5 region proteins can be produced synthetically by chemical reaction or by recombinant techniques in conjunction with expression systems. These deglycosylated kringle 1-5 region proteins can be animal or human in origin. In one embodiment, the deglycosylated kringle 1-5 region protein is produced recombinantly and has the amino acid sequence of SEQ ID NO:1.

Deglycosylated kringle 1-5 region proteins may also be produced *in vitro* or *in vivo* by enzymatically cleaving plasminogen or plasmin to generate proteins having antiangiogenic activity or by using compounds that mimic the action of endogenous enzymes that cleave deglycosylated plasminogen into deglycosylated kringle 1-5 region proteins. Deglycosylated kringle 1-5 region protein production may also be modulated by compounds that affect the activity of plasminogen cleaving enzymes. In particular, it is a surprising discovery of the present invention that glycosylation of plasminogen reduces the yield of kringle 1-5 region proteins upon cleavage of plasminogen with elastase. Therefore, in one

embodiment of the present invention, the method of making deglycosylated kringle 1-5 region proteins comprises cleaving deglycosylated plasminogen with elastase. Given the role of elastase reported here in the generation of kringle 1-5 region proteins, it is likely that the N-linked carbohydrate on kringle 3 will also modulate the reaction mechanism of other proteinases potentially responsible for kringle 1-5 region protein generation.

The present invention also includes methods for the detection of deglycosylated kringle 1-5 region proteins in body fluids and tissues for the purpose of determining the efficacy of treatment and for the prognosis and diagnosis of angiogenesis associated diseases such as cancer. In particular, provided herein are methods of using antibodies that are specific for deglycosylated kringle 1-5 region proteins over circulating plasminogen. These methods provide a means to detect and distinguish deglycosylated kringle 1-5 region proteins from glycosylated kringle 1-5 region proteins and glycosylated plasminogen. The present invention also includes the detection of deglycosylated kringle 1-5 region protein binding sites and receptors in cells and tissues.

Kits for detection and measurement of deglycosylated kringle 1-5 region proteins, and the receptors therefor, are contemplated as part of the present invention. Antisera that possess the highest titer and specificity and can detect deglycosylated kringle 1-5 region proteins in extracts of plasma, urine, tissues, and in cell culture media are further examined to establish easy to use kits for rapid, reliable, sensitive, and specific measurement and localization of deglycosylated kringle 1-5 region proteins. These assay kits include but are not limited to the following techniques; competitive and non-competitive assays, radioimmunoassay, bioluminescence and chemiluminescence assays, fluorometric assays, sandwich assays, immunoradiometric assays, dot blots, enzyme linked assays including ELISA, antibody coated strips or dipsticks for rapid monitoring of urine or blood, and immunocytochemistry. For

each kit the range, sensitivity, precision, reliability, specificity and reproducibility of the assay are established. Intra-assay and inter-assay variation is established at 20%, 50% and 80% points on the standard curves of displacement or activity.

5 The present invention also includes methods of treating or preventing angiogenic diseases and processes including, but not limited to, arthritis and tumors by stimulating the production of deglycosylated kringle 1-5 region proteins, and/or by administering substantially purified deglycosylated kringle 1-5 region proteins, nucleotides encoding deglycosylated kringle 1-5 region proteins, or deglycosylated kringle 1-5 region protein agonists or antagonists, and/or deglycosylated kringle 1-5 region protein antisera or antisera directed against deglycosylated kringle 1-5 region protein antisera to a patient. These angiostatin deglycosylated kringle 1-5 region proteins, deglycosylated kringle 1-5 region protein antisera, deglycosylated kringle 1-5 region protein receptor agonists or antagonists, or combinations thereof, are combined with pharmaceutically acceptable excipients, and optionally a sustained-release matrix, such as biodegradable polymers, to form therapeutic compositions. Additional treatment methods include administration of deglycosylated kringle 1-5 proteins, deglycosylated kringle 1-5 region protein analogs, deglycosylated kringle 1-5 region protein antisera, or deglycosylated kringle 1-5 region protein receptor agonists and antagonists linked to cytotoxic agents.

25 In one aspect of the present invention, a method of inhibiting angiogenesis in an individual comprises, increasing in the individual an *in vivo* concentration of a deglycosylated fragment of a kringle 1-5 region of a plasminogen protein relative to the *in vivo* concentrations of a glycosylated fragment of a kringle 1-5 region of a plasminogen protein and wherein the deglycosylated fragment of the kringle 1-5 region protein has antiangiogenic activity *in vivo*. The *in vivo* concentration of the deglycosylated fragment can be increased in any bodily fluid or tissue, but is preferably increased in the serum. In a further

embodiment, a deglycosylated kringle 1-5 region protein is administered in a treatment effective amount to an individual in need of such treatment. In another embodiment, a gene encoding a deglycosylated fragment of a kringle 1-5 region of a plasminogen is administered to an individual using the gene therapy methods discussed in more detail below. Preferably the gene contains an amino acid substitution that prevents glycosylation during post-translational modification of the fragment of the kringle 1-5 region, and more preferably the substitution is at an amino acid corresponding to amino acid Asn-289 of human plasminogen.

A deglycosylated kringle 1-5 region protein is effective in treating diseases or processes that are mediated by, or involve, angiogenesis. The present invention includes the method of treating an angiogenesis mediated disease with an effective amount of a deglycosylated kringle 1-5 region protein, or combinations of deglycosylated kringle 1-5 region proteins that collectively possess antiangiogenic activity, or deglycosylated kringle 1-5 region protein agonists and antagonists. The angiogenesis mediated diseases include, but are not limited to, solid tumors; blood born tumors such as leukemias; tumor metastasis; benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; rheumatoid arthritis; psoriasis; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; and wound granulation. Deglycosylated kringle 1-5 region protein is useful in the treatment of disease of excessive or abnormal stimulation of endothelial cells. These diseases include, but are not limited to, intestinal adhesions, Crohn's disease, atherosclerosis, scleroderma, and hypertrophic scars, i.e., keloids. Deglycosylated kringle 1-5 region proteins can be used as a birth

control agent by preventing vascularization required for embryo implantation. Deglycosylated kringle 1-5 region proteins are also useful in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochela minialia quintosa*) and ulcers (*Helicobacter pylori*).

Deglycosylated kringle 1-5 region proteins may be used in combination with other compositions and procedures for the treatment of diseases. For example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with deglycosylated kringle 1-5 region proteins and then deglycosylated kringle 1-5 region proteins may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize and inhibit the growth of any residual primary tumor.

In addition to methods of administering deglycosylated kringle 1-5 region proteins, included in the present invention are methods of passive antibody therapy. In one embodiment, deglycosylated kringle 1-5 region protein antibodies block the action of excess endogenous deglycosylated kringle 1-5 region proteins. Specifically blocking the action of excess endogenous deglycosylated kringle 1-5 region proteins (versus endogenous glycosylated kringle 1-5 region proteins) is important considering the novel teaching herein that deglycosylated kringle 1-5 region proteins are more antiangiogenic than glycosylated kringle 1-5 region proteins. This treatment provides an improved method of modulating angiogenic-dependent processes such as reproduction, development, and wound healing and tissue repair and treating abnormal ovulation, menstruation and placentation, and vasculogenesis. This method also provides a useful tool to examine the effects of deglycosylated kringle 1-5 region protein removal on metastatic processes.

In another embodiment, antisera directed to the Fab regions of deglycosylated kringle 1-5 region protein antibodies can be administered to block the ability of endogenous deglycosylated kringle 1-5 region protein antisera to bind

deglycosylated kringle 1-5 region proteins. The net effect of this treatment is to facilitate the ability of endogenous circulating deglycosylated kringle 1-5 region protein to reach target cells, thereby decreasing the spread of metastases.

5 The deglycosylated kringle 1-5 region proteins and antibodies thereto described above can be provided as isolated and substantially purified proteins and protein fragments in pharmaceutically acceptable formulations using formulation methods known to those of ordinary skill in the art. The
10 deglycosylated kringle 1-5 region proteins and antibodies thereto described above may be a solid, liquid or aerosol. Examples of solid therapeutic compositions include pills, creams, and implantable dosage units. The pills may be administered orally and the therapeutic creams may be administered topically. The
15 implantable dosage units may be administered locally, for example at a tumor site, or may be implanted for systemic release of the therapeutic angiogenesis-modulating composition, for example subcutaneously. Examples of liquid compositions include formulations adapted for injection subcutaneously,
20 intravenously, intraarterially, and formulations for topical and intraocular administration. Examples of aerosol formulations include inhaler formulations for administration to the lungs. In general however, the formulations may be administered by any route, including but not limited to, the topical, transdermal,
25 intraperitoneal, intracranial, intracerebroventricular, intracerebral, intravaginal, intrauterine, oral, rectal or parenteral (e.g., intravenous, intraspinal, subcutaneous or intramuscular) route.

In addition, the deglycosylated kringle 1-5 region proteins and antibodies thereto may be incorporated into biodegradable
30 polymers allowing for sustained release of the compound, the polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the deglycosylated kringle 1-5 region protein or antibody is slowly released systemically. The biodegradable polymers and
35 their use are described, for example, in detail in Brem *et al.*,

“Interstitial chemotherapy with drug polymer implants for the treatment of recurrent gliomas” *J. Neurosurg.* 74:441-446 (1991). Osmotic minipumps may also be used to provide controlled delivery of high concentrations of deglycosylated kringle 1-5 region proteins and antibodies thereto through cannulae to the site of interest, such as directly into a metastatic growth or into the vascular supply to that tumor.

The dosage of the deglycosylated kringle 1-5 region proteins and antibodies of the present invention will depend on the disease state or condition being treated and other clinical factors such as weight and condition of the human or animal and the route of administration of the compound. For treating humans or animals, between approximately 0.5 mg/kilogram per day to 500 mg/kilogram per day of the deglycosylated kringle 1-5 region protein can be administered. Depending upon the half-life of the deglycosylated kringle 1-5 region protein in the particular animal or human, it can be administered between several times per day to once a week. It is to be understood that the present invention has application for both human and veterinary use. The methods of the present invention contemplate single as well as multiple administrations, given either simultaneously or over an extended period of time.

The deglycosylated kringle 1-5 region protein formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). Suitable pharmaceutical carriers and excipients are known to those skilled in the art, however, an example of a suitable pharmaceutical excipient is water. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

5 Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules or vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

10 Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients, particularly mentioned above, the formulations of the present invention may include other agents conventional in the art having regard to the type of formulation in question. Optionally, cytotoxic agents may be incorporated or otherwise combined with deglycosylated kringle 1-5 region proteins, or biologically functional protein fragments thereof, to provide dual therapy to the patient.

15 The present invention also encompasses gene therapy whereby the gene encoding a deglycosylated kringle 1-5 region protein is regulated in a patient. Various methods of transferring or delivering DNA to cells for expression of the gene product, otherwise referred to as gene therapy, are disclosed in "Gene Transfer into Mammalian Somatic Cells *in vivo*", N. Yang, *Crit. Rev. Biotechn.* 12(4):335-356 (1992). Gene therapy encompasses incorporation of DNA sequences into somatic cells or germ line cells for use in either *ex vivo* or *in vivo* therapy. Gene therapy functions to replace genes, augment normal or abnormal gene function, and to combat infectious diseases and other pathologies.

5 Strategies for treating medical problems with gene therapy include therapeutic strategies such as replacing a defective gene with a functional gene or adding a functional gene to augment a slightly functional gene. The genes inserted may treat a disease or condition or make the tissue or organ more susceptible to a treatment regimen. With regard to the present invention, a gene for a deglycosylated kringle 1-5 region protein may be placed in a subset of cells, thus preventing the occurrence of angiogenesis in the transformed cells.

10 Many protocols for the transfer of deglycosylated kringle 1-5 region protein DNA or deglycosylated kringle 1-5 region protein regulatory sequences are envisioned in this invention. Transfection of promoter sequences, other than one normally found specifically associated with a deglycosylated kringle 1-5 region protein, or other sequences that would increase production of deglycosylated kringle 1-5 region proteins are envisioned as methods of gene therapy. Such "genetic switches" could be used to activate a deglycosylated kringle 1-5 region protein (or the deglycosylated kringle 1-5 region protein receptor) in cells not normally expressing deglycosylated kringle 1-5 region protein (or the deglycosylated kringle 1-5 region protein receptor).

25 Gene transfer methods for gene therapy fall into three broad categories: (1) chemical (lipid-based carriers, or other non-viral vectors), (2) biological (virus-derived vector and receptor uptake), and (3) physical (electroporation, direct gene transfer and particle bombardment). Gene therapy methodologies can also be described by delivery site. Fundamental ways to deliver genes include *ex vivo* gene transfer, *in vitro* gene transfer, and *in vivo* gene transfer. In *ex vivo* gene transfer, cells are taken from the patient and grown in cell culture. The DNA is transfected into the cells, the transfected cells are expanded in number and then re-implanted in the patient. The present invention encompasses the removal of endothelial cells from a patient, transfection of DNA encoding a deglycosylated kringle 1-5

region protein, or regulatory sequence thereof, and re-introduction of the transfected endothelial cells into the patient. In *in vitro* gene transfer, transformed cells, such as endothelial cells, growing in culture are introduced into the patient. The transformed cells are not taken from the patient who will receive the gene therapy. *In vivo* gene transfer involves introducing the DNA into the cells of the patient when the cells are within the patient. Methods include using a noninfectious virus to introduce a gene into a patient or injecting naked DNA into a site in the patient whereby DNA is taken up by a percentage of cells in which the gene product protein is expressed. In the present invention, DNA encoding a deglycosylated kringle 1-5 region protein can be introduced into the endothelial cells lining the blood vessels, thereby inhibiting angiogenesis. In a preferred embodiment, the DNA encoding a deglycosylated kringle 1-5 region protein is introduced into endothelial cells lining the blood vessels in close proximity to a tumor.

Chemical methods of gene therapy may involve a lipid based compound, not necessarily a liposome, used to ferry the DNA across the cell membrane. Lipofectins or cytofectins, lipid-based positive ions that bind to negatively charged DNA, make a complex that can cross the cell membrane and provide the DNA into the interior of the cell. Liposome/DNA complexes may be directly injected intravenously into the patient. It is believed that the liposome/DNA complexes are concentrated in the liver where they deliver the DNA to macrophages and Kupffer cells. These cells are long lived and thus provide long term expression of the delivered DNA. Additionally, vectors or the "naked" DNA of the gene may be directly injected into the desired organ, tissue or tumor for targeted delivery of the therapeutic DNA. Other DNA carrier systems include the asialoglycoprotein/polylysine conjugate system for carrying DNA to hepatocytes for *in vivo* gene transfer and DNA coupled to nuclear proteins in specifically engineered vesicle complexes that are carried directly into the nucleus.

Biological methods used in gene therapy techniques may involve receptor-based endocytosis, or receptor-based phagocytosis, which involve binding a specific ligand to a cell surface receptor and enveloping and transporting the ligand across the cell membrane. Specifically, a ligand/gene complex is created and injected into the blood stream. Target cells having a receptor for the ligand will specifically bind the ligand and transport the ligand-DNA complex into the cell. Additional biological methods employ viral vectors to insert genes into cells. For example, altered retrovirus vectors have been used in *ex vivo* methods to introduce genes into peripheral and tumor-infiltrating lymphocytes, hepatocytes, epidermal cells, myocytes, and other somatic cells. These altered cells are then introduced into the patient.

Viral vectors have also been used to insert genes into cells using *in vivo* protocols. To accomplish tissue-specific expression of foreign genes, cis-acting regulatory elements or promoters that are known to be tissue specific can be used. Alternatively, tissue-specific expression can be achieved using *in situ* delivery of DNA or viral vectors to specific anatomical sites *in vivo*. For example, gene transfer to blood vessels *in vivo* has been achieved by implanting *in vitro* transduced endothelial cells in chosen sites on arterial walls. Surrounding cells were infected by the virus and therefore also expressed the gene product. A viral vector can be delivered directly to the *in vivo* site, by a catheter for example, thus allowing only certain areas to be infected by the virus. *In vivo* gene transfer using retrovirus vectors has also been demonstrated in mammary tissue and hepatic tissue by injection of the altered virus into blood vessels leading to the organs.

Viral vectors that have been used for gene therapy protocols include, but are not limited to, retroviruses such as murine leukemia retroviruses, RNA viruses such as poliovirus or Sindbis virus, adenovirus, adeno-associated virus, herpes viruses, SV40, vaccinia and other DNA viruses. Replication-defective

murine retroviral vectors are the most widely utilized gene transfer vectors. Fundamental advantages of retroviral vectors for gene transfer include efficient infection and gene expression in most cell types, precise single copy vector integration into target cell chromosomal DNA, and ease of manipulation of the retroviral genome. The adenovirus is capable of transducing novel genetic sequences into target cells *in vivo*. Adenoviral-based vectors express gene product proteins at high levels and have high efficiencies of infectivity, even with low titers of virus. Additionally, the virus is fully infective as a cell free virion so injection of expression cell lines is not necessary. Another potential advantage to adenoviral vectors is the ability to achieve long term expression of heterologous genes *in vivo*.

Mechanical methods of DNA delivery include direct injection of DNA, such as microinjection of DNA into germ or somatic cells, pneumatically delivered DNA-coated particles, such as the gold particles used in a "gene gun," inorganic chemical approaches such as calcium phosphate transfection and electroporation. It has been found that injecting plasmid DNA into muscle cells yields high percentage of the cells that are transfected and have sustained expression of marker genes. The DNA of the plasmid may or may not integrate into the genome of the cells. Non-integration of the transfected DNA would allow the transfection and expression of gene product proteins in terminally differentiated, non-proliferative tissues for a prolonged period of time without fear of mutational insertions, deletions, or alterations in the cellular or mitochondrial genome. Long-term, but not necessarily permanent, transfer of therapeutic genes into specific cells may provide treatments for genetic diseases or for prophylactic use. The DNA could be re-injected periodically to maintain the gene product level without mutations occurring in the genomes of the recipient cells. Non-integration of exogenous DNAs may allow for the presence of several different exogenous DNA constructs within one cell with all of the constructs expressing various gene products.

Both particle-mediated gene transfer methods and electroporation can be used in *in vitro* systems, or with *ex vivo* or *in vivo* techniques to introduce DNA into cells, tissues or organs. With regard to particle-mediated gene transfer, a particle bombardment device, or "gene gun," is used that generates a motive force to accelerate DNA-coated high density particles (such as gold or tungsten). These particles penetrate the target organs, tissues or cells. Electroporation mediated gene transfer comprises the use of a brief electric impulse with a given field strength that is used to increase the permeability of a membrane in such a way that DNA molecules can penetrate into the cells.

The gene therapy protocol for transfecting DNA encoding deglycosylated kringle 1-5 region proteins into a patient may either be through integration of the deglycosylated kringle 1-5 region protein DNA into the genome of the cells, into minichromosomes or as a separate replicating or non-replicating DNA construct in the cytoplasm or nucleoplasm of the cell. Deglycosylated kringle 1-5 region protein expression may continue for a long-period of time or the DNA may be re-injected periodically to maintain a desired level of the deglycosylated kringle 1-5 region protein in serum or in a cell, tissue or organ.

This invention is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon the scope thereof. It is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention and/or the scope of the appended claims.

Example 1

Isolation of glycosylated and deglycosylated kringle 1-5 region proteins

Human plasminogen was purified using affinity chromatography as described in Brockway, W.J. and Castellino, F.J., "Measurement and the binding of antifibrinolytic amino acids to various plasminogens" *Arch. Biochem. Biophys.* 151:194-199 (1972). One liter of human plasma (Children's Hospital blood bank) was applied to a 200mL lys-Sepharose 4b (Pharmacia) column at a flow rate of 2-3 ml/minute. The column, previously equilibrated with 50mM Tris pH 7.4 (Sigma) was then washed with 500mL 50mM tris/1M NaCl pH 7.4. Bound plasminogen was eluted with 200mL 50mM Tris/200mM ϵ -aminocaproic acid (ϵ ACA). The purity of this material was greater than 95% as assessed by SDS-PAGE. Human plasminogen was further fractionated into glycosylated plasminogen (plasminogen 1) and deglycosylated plasminogen (plasminogen 2) using lectin affinity chromatography. Human plasminogen (10mg) was applied to a 5mL conA HiTrap column (Pharmacia), equilibrated with 50mM Tris pH 7.4/1mM MgCl_2 /1mM CaCl_2 . Plasminogen 2, lacking an N-linked carbohydrate, does not bind to this column. After washing the column with five column volumes of equilibration buffer, plasminogen 1 was eluted using a solution of 50mM Tris pH 7.4/1mM MgCl_2 /1mM CaCl_2 . Protein concentration was determined spectrophotometrically as a wavelength of 280nm using an $A^{0.1\%/1\text{cm}}$ of 1.6 (computed from the protparam tool available at <http://www.expasy.ch/>).

Plasminogen 1 and 2 were dialyzed against 20mM Tris pH 7.6. Equal amounts of these glycoforms (as determined by A280nm) were then digested with porcine pancreatic elastase (PPE) as described in O'Reilly, M.S. *et al.*, "Angiostatin induces and sustains dormancy of human primary tumors in mice" *Nature Medicine* 2:689-692 (1996). Essentially, plasminogen was digested at 37°C with 0.8 units of PPE (Calbiochem) per

milligram of plasminogen. After five hours incubation, the reaction was quenched by applying the reaction mixture to a lys-Sepharose 4b column (10mL), and eluting an approximately kringle 1-3 fragment of plasminogen and other lysine binding fragments, such as kringle 4, with 5mL 200mM εACA/50mM Tris pH 7.4. The kringle 1-3 region protein was separated from kringle 4 and other smaller (<12kDa) fragments using gel filtration on a Superdex200 HR 10/30 column (Pharmacia). The Superdex200 column was equilibrated with PBS. Kringle 1-3 region protein concentration was determined spectrophotometrically at a wavelength of 280nm, using $A^{0.1\%/1\text{cm}}$ value of 1.87 (computed from the protparam tool available at <http://www.expasy.ch/>).

Bovine capillary endothelial cells were obtained as described previously in O'Reilly, M.S. *et al.* "Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma" *Cell* 79:315-328 (1994) and were maintained in DMEM with 10% heat-inactivated BCS, antibiotics, and 3ng/mL recombinant human bFGF (Scios Nova, CA). Cells were washed with PBS and dispersed in a 0.05 % solution of trypsin. A cell suspension was made with DMEM/10% BCS/1% antibiotics and the concentration adjusted to 25,000 cells/mL. Cells were plated onto gelatinized 24-well culture plates (0.5mL/well) and were incubated at 37°C in 10% CO₂ for 24 hours. The media was replaced with 0.25mL of DMEM/5% BCS/1% antibiotics, and the test sample applied. After 20 minutes incubation, media and bFGF were added to each well to obtain a final volume of 0.5mL DMEM/5% BCS/1% antibiotics/1 ng/mL bFGF. After 72 hours, cells were dispersed in trypsin, resuspended in Hematell (Fisher scientific, PA) and counted by Coulter counter. Data from endothelial cell proliferation assays were plotted and IC₅₀ values determined using SigmaPlot. All data points were determined in triplicate from four separate preparations of kringle 1-5 region protein.

Example 2

Deglycosylation of plasminogen results in an increased yield of kringle 1-5 region proteins

Equal amounts of plasminogen 1 and plasminogen 2 were digested with PPE, and the kringle 1-5 region proteins purified as described in Example 1 using a combination of affinity chromatography (lys-Sepharose) to purify kringle 1-5 region proteins and other lysine binding fragments of plasminogen, and gel filtration (Superdex200) to separate kringle 1-5 region proteins from kringle 4. Figure 2 shows a typical chromatogram from the affinity column, demonstrating the reduced yield obtained when plasminogen 1 is used, as a substrate compared to plasminogen 2. Figure 3 shows the chromatogram obtained from a typical gel filtration experiment as described in Example 1). It was noted that kringle 1-5 region protein from plasminogen 1 has a slightly lower retention time (13.77 minutes) than deglycosylated kringle 1-5 region protein (14.36 minutes), as would be expected from this glycosylated fragment. The yield of kringle 1-5 region proteins from plasminogen 2 was 4-5 fold higher than the yield of kringle 1-5 region proteins using plasminogen 1 as a substrate. Glycosylated kringle 1-5 region proteins migrated at approximately 40 kDa and deglycosylated kringle 1-5 region proteins migrated at 38kDa on SDS-PAGE (data not shown). Edman degradation amino-terminal sequence analysis of both kringle 1-5 region protein glycoforms, gave identical sequence starting at Arg87, 16 residues amino-terminal to the cysteine residue that marks the beginning of kringle 1.

Therefore, it was determined that the presence of N-linked carbohydrate on Asn-289 modulates the generation of kringle 1-5 region proteins from human plasminogen. The presence of carbohydrate results in a 4-5-fold lower yield of kringle 1-5 region protein when PPE is used to cleave kringle 1-5 region proteins out of plasminogen. It was also determined that fractionation of kringle 1-5 region proteins generated from material containing glycosylated plasminogen (plasminogen 1)

and deglycosylated plasminogen (plasminogen 2) in the ratio 40:60 using lectin affinity chromatography resulted in only 5-10% of glycosylated kringle 1-5 region proteins (data not shown). This data further indicated that N-linked carbohydrate is a negative modulator of kringle 1-5 region protein generation. Both kringle 1-5 region glycoforms had identical N-terminal sequence, indicating that the carbohydrate did not interfere with the site of proteolytic cleavage.

Although glycosylated kringle 1-5 region proteins were less efficiently generated and less efficient as an inhibitor of angiogenesis, it may be that kringle 1-5 region proteins containing N-linked carbohydrate, will have a longer clearance time than deglycosylated kringle 1-5 region proteins. This may be a mechanism to ensure persistence of anti-endothelial activity systemically, albeit at a reduced level.

Example 3

Deglycosylated kringle 1-5 region proteins have increased antiangiogenic activity as compared to glycosylated kringle 1-5 region proteins

Both kringle 1-5 region protein glycoforms were assayed for their ability to inhibit the proliferation of bovine endothelial cells as described in Example 1. Both kringle 1-5 region protein glycoforms inhibited endothelial cell proliferation. However, glycosylated kringle 1-5 region protein, which contains an N-linked carbohydrate, has a measured IC_{50} value of approximately $13\mu g/mL$, whereas deglycosylated kringle 1-5 region protein, the glycoform, lacking an N-linked carbohydrate, has a measured IC_{50} value of approximately $2.4\mu g/mL$. Thus, deglycosylated kringle 1-5 region protein appears to be a much more efficient inhibitor of endothelial cell proliferation, and in particular was 4-5 fold more efficient as an inhibitor of endothelial cell proliferation.

Claims

We claim:

1. A composition comprising a pharmaceutically acceptable carrier and a deglycosylated fragment of a kringle 1-5 region of a plasminogen protein in a greater amount than a naturally glycosylated form of the deglycosylated fragment, wherein the deglycosylated fragment lacks one or more carbohydrate moieties linked to the naturally glycosylated form and wherein the deglycosylated fragment has antiangiogenic activity.

2. The composition of claim 1, wherein the deglycosylated fragment lacks a bisialylated-biantennary glycan.

3. The composition of claim 1, wherein the deglycosylated fragment lacks an N-linked carbohydrate moiety.

4. The composition of claim 1, wherein the deglycosylated fragment lacks a carbohydrate chain at amino acid position Asn-289 of human plasminogen.

5. The composition of claim 1, wherein the deglycosylated fragment is approximately a kringle 1-3 protein.

6. The composition of claim 5, wherein the deglycosylated fragment begins at approximately amino acid 87 of human plasminogen.

7. The composition of claim 5, wherein the deglycosylated fragment amino acid sequence is shown in SEQ ID NO:1.

8. The composition of claim 1, wherein the deglycosylated fragment is produced recombinantly.

9. The composition of claim 1, wherein the deglycosylated fragment has an amino acid substitution at amino acid position 289 of human plasminogen.

10. The composition of claim 1, wherein the deglycosylated fragment and the glycosylated form are at a ratio of at least 60:40.

11. The composition of claim 1, wherein the deglycosylated fragment and the glycosylated form are at a ratio of at least 80:20.

12. The composition of claim 1, wherein the deglycosylated fragment and the glycosylated form are at a ratio of 100:0.

13. A nucleotide sequence encoding the deglycosylated fragment of claim 1.

14. The nucleotide sequence of claim 13, wherein the nucleotide sequence is shown in SEQ ID NO:2.

15. The composition of claim 1, wherein the deglycosylated fragment has antiangiogenic activity *in vitro*.

16. The composition of claim 1, wherein the deglycosylated fragment has antiangiogenic activity *in vivo*.

17. A method of inhibiting angiogenesis in an individual comprising, increasing in the individual an *in vivo* concentration of a deglycosylated fragment of a kringle 1-5 region of a plasminogen protein relative to *in vivo* concentration of a naturally glycosylated form of the deglycosylated fragment, wherein the deglycosylated fragment lacks one or more

carbohydrate moieties linked to the naturally glycosylated form and wherein the deglycosylated fragment has antiangiogenic activity *in vivo*.

5 18. The method of claim 17, wherein the deglycosylated fragment is administered to the individual.

10 19. The method of claim 17, wherein the deglycosylated fragment lacks a bisialylated-biantennary glycan.

10 20. The method claim 19, wherein the deglycosylated fragment lacks an N-linked carbohydrate.

15 21. The method of claim 20, wherein the deglycosylated fragment lacks a carbohydrate chain at amino acid position Asn-289 of human plasminogen.

20 22. The method of claim 17, wherein the deglycosylated fragment is approximately a kringle 1-3 protein.

20 23. The method of claim 22, wherein the deglycosylated fragment begins at approximately amino acid 87 of human plasminogen.

25 24. The method of claim 22, wherein the deglycosylated fragment amino acid sequence is shown in SEQ ID NO:1.

30 25. The method of claim 17, wherein the deglycosylated fragment is produced recombinantly.

35 26. The method of claim 17, wherein the deglycosylated fragment has an amino acid substitution at amino acid position 289 of human plasminogen.

27. A deglycosylated fragment of a kringle 1-5 region of a plasminogen protein, wherein the deglycosylated fragment amino acid sequence is shown in SEQ ID NO:1.

5

28. A nucleic acid encoding a deglycosylated fragment of a kringle 1-5 region of a plasminogen protein, wherein the deglycosylated fragment nucleic acid sequence is shown in SEQ ID NO:2.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
221

**DEGLYCOSYLATED KRINGLE 1-5 REGION
FRAGMENTS OF PLASMINOGEN AND
METHODS OF USE**

5

Abstract of the Invention

10 Disclosed are deglycosylated fragments of a kringle 1-5
region of plasminogen, nucleotides encoding deglycosylated
kringle 1-5 region proteins and antibodies specific for
deglycosylated kringle 1-5 region proteins. The compositions of
the present invention have increased antiangiogenic activity as
15 compared to previously isolated kringle 1-5 region proteins.
Also included in the present invention are methods of treating
angiogenesis-associated diseases and conditions such as cancer
using the compositions described herein.

Attorney Docket: 05940-0141

SEQ ID NO:1
 VYLSECKTGNGKNYRGTMSTKNGITCQKWSSTSPHRPR
 FSPATHPSEGLEENYCRNPDNDPQGPWCYTTDPEKRYDY
 5 CDILECEEECMHCSGENYDGGKISKTMSGLECQAWDSQSP
 HAHGYPISKFPNKNLKNYCRNPDRELRPWCFTTDPNKR
 WELCDIPRCTTPPPSSGPTYQCLKGTGENYRGNVAVTVSG
 HTCQHWSAQTPHHERTPENFPCKNLDENYCRNPDGKR
 APWCHTTNSQVRWEYCKIPSCDSSPV

SEQ ID NO:2
 GTGTATCTCTCAGAGTGCAAGACTGGGAATGGAAAGAA
 TTACAGAGGGACGATGTCCAAAAACAAAAATGGCATC
 15 ACCTGTCAAAAAATGGAGTTCCACTTCTCCCCACAGACCT
 AGATTCTCACCTGCTACACACCCCTCAGAGGGACTGGA
 GGAGAACTACTGCAGGAATCCAGACAACGATCCGCAG
 GGGCCCTGGTGCTATACTACTGATCCAGAAAAGAGATA
 TGACTACTGCGACATTCTTGAGTGTGAAGAGGAATGTA
 TGCATTGCAGTGGAGAAAATATGACGGCAAAATTTCC
 20 AAGACCATGTCTGGACTGGAATGCCAGGCCTGGGACTC
 TCAGAGCCCACACGCTCATGGATACATTCCCTTCAAATT
 TCCAAACAAGAACCTGAAGAAGAATTACTGTGCTAACC
 CCGATAGGGAGCTGCGGCCTTGGTGTTTACCACCCGAC
 CCAACAAGCGCTGGGAACCTTGTGACATCCCCCGCTG
 25 CACAACACCTCCACCATCTTCTGGTCCACCTACCACTG
 TCTGAAGGGAACAGGTGAAAATATCGCGGGAATGTG
 GCTGTTACCGTGTCCGGGCACACCTGTCAGCACTGGAGT
 GCACAGACCCCTCACACACATGAAAGGACACCAGAAA
 ACTTCCCCTGCAAAAAATTGGATGAAAATCTGCCGC
 30 AATCCTGACGGAAAAAGGGCCCCATGGTGCCATACAAC
 CAACAGCCAAGTGCAGTGGGAGTACTGTAAGATACCGT
 CCTGTGACTCCTCCCCAGTA

Figure 1




Figure 2




Figure 3




Figure 4